arXiv:1404.6009v1 [math.RA] 24 Apr 2014

EXPLICIT IDEMPOTENTS OF FINITE GROUP ALGEBRAS

F. E. BROCHERO MARTÍNEZ AND C. R. GIRALDO VERGARA

Abstract. Let $\mathbb{F}_q$ be a finite field, G a finite cyclic group of order p^k and p is an odd prime with $\gcd(q,p)=1$. In this article, we determine an explicit expression for the primitive idempotents of $\mathbb{F}_qG$. This result extends the result in [1], [2] and [8].

1. Introduction

Let G be a finite cyclic group of order n and $\mathbb{F}_q$ a finite field of order q, where q is prime relative to n. The cyclic codes of length n over $\mathbb{F}_q$ can be viewed as an ideal in the group algebra $\mathbb{F}_qG$ and each ideal is generated by an idempotent of $\mathbb{F}_qG$. By the representation theorem of abelian groups we know that

$$G \simeq C_{p_1^{\beta_1}} \times \cdots \times C_{p_r^{\beta_r}}$$

where $C_{p_j^{\beta_j}}$ is a cyclic group of order $p_j^{\beta_j}$ and $p_1, \ldots, p_r$ are distinct primes. In addition, it is well known that

$$\mathbb{F}_qG \simeq \mathbb{F}_qC_{p_1^{\beta_1}} \otimes \cdots \otimes \mathbb{F}_qC_{p_r^{\beta_r}}.$$

From this fact, in order to construct the idempotents of the cyclic group algebra $\mathbb{F}_qG$, it is enough to consider the case $G = C_n$ where n is a power of a prime. Observe that the condition $\gcd(n,q)=1$ is necessary by the Maschke theorem (see [4] theorem 10.8).

2. Primitive idempotents: General calculation

Let $\Phi_d(x)$ denote the d-th cyclotomic polynomial, i.e., $\Phi_d(x)$ can be defined recursively by $\Phi_1(x) = x-1$ and $x^k - 1 = \prod_{d|k} \Phi_d(x)$. It is well known (see [5] page 65 theorem 2.47) that if $\gcd(q,d)=1$ then $\Phi_d(x)$ can be factorized into $r_d = \frac{\varphi(d)}{s_d}$ distinct monic irreducible polynomials of the same degree s_d over $\mathbb{F}_q$ and $s_d = \operatorname{ord}_d q = \min\{k \in \mathbb{N}^* | q^k \equiv 1 \pmod d\}$, i.e. $\Phi_d(x)$ can be factorized in $\mathbb{F}_q[x]$ as $f_{d,1} \cdot f_{d,2} \cdots f_{d,r_d}$, where each $f_{d,j}$ is an irreducible polynomial of degree s_d, and then

$$x^n - 1 = \prod_{d|n} \prod_{j=1}^{r_d} f_{s,j}.$$

Observe that if K is a decomposition field of the cyclotomic polynomial $\Phi_d(x)$, then for each pair $f_{d,i}, f_{d,j}$ there exists $\tau \in Gal(K|\mathbb{F}_q)$ such that $\tau(f_{d,i}) = f_{d,j}$.

Date: April 25, 2014.
2010 *Mathematics Subject Classification.* 16S34(primary) and 94B05(secondary).
Key words and phrases. Irreducible cyclic codes, Primitive Idempotents.

By the Chinese remainder theorem, we know that

$$\mathbb{F}_q C_n \simeq \frac{\mathbb{F}_q[x]}{\langle x^n-1\rangle} \simeq \bigoplus_{d|n}\bigoplus_{j=1}^{r_d} \frac{\mathbb{F}_q[x]}{\langle f_{d,j}\rangle}$$

where the $\mathbb{F}_q$-algebra isomorphisms are naturally defined using a generator g of C_n as $g \mapsto \overline{x} \mapsto (\overline{x},\dots,\overline{x})$.

Since each direct sum term is a field, then this decomposition is a Weddeburn decomposition of the group algebra and each primitive idempotent is of the form $(\overline{0},\dots,\overline{0},\overline{1},\overline{0},\dots,\overline{0})$. Therefore, if $e_{d,j}$ is a primitive idempotent of $\mathbb{F}_q C_n$, then it can be seen as a polynomial $e_{d,j}(x)$ with the following properties :

(1) $deg(e_{d,j}(x)) < n$
(2) $e_{d,j}(x)$ is divisible by f_{d_1,j_1} for all $(d_1,j_1) \neq (d,j)$
(3) $e_{d,j}(x) - 1$ is divisible by $f_{d,j}$.

From these three properties, we have the following

Theorem 2.1. *Let $\mathbb{F}_q$ be a finite field with q element and $n \in \mathbb{N}^*$ such that $gcd(q,n) = 1$, then each primitive idempotent of $\mathbb{F}_q C_n$ is of the form*

$$e_{d,j}(x) = \frac{x^n-1}{f_{d,j}(x)} h_{d,j}(x),$$

where $f_{d,j}$ is an irreducible factor of the cyclotomic polynomial $\Phi_d(x)$, d is a divisor of n and $h_{d,j} \in \mathbb{F}_q[x]$ is a polynomial with $\deg(h_{d,j}) < s_d := \operatorname{ord}_d q$ that is the inverse of $\frac{x^n-1}{f_{d,j}(x)}$ in the field $\frac{\mathbb{F}_q[x]}{\langle f_{d,j}\rangle}$.

Observe that if we know the polynomial $f_{d,j}$ then $h_{d,j}$ can be explicitly calculated using the Extended Euclidean Algorithm for polynomials. In general, the factorization of $\Phi_d(x)$ in $\mathbb{F}_q[x]$ for arbitrary d and q is an open problem. Some especial cases can be found in [5], [6] and [7].

3. The idempotents for some special known cases

In this section we are going to show, without proof, the idempotents in the extremal cases where $x^n - 1$ factorized into linear factors in $\mathbb{F}_q[x]$ and where each cyclotomic polynomial $\Phi_d(x)$, with $d|n$, is irreducible in $\mathbb{F}_q[x]$. Before that, we need the following remarks:

Remark 3.1. *The cyclotomic polynomial $\Phi_d(x) \in \mathbb{F}_q[x]$, with $\gcd(d,q) = 1$ is factorized into linear factors if and only if $q \equiv 1 \pmod d$, thus, $x^n - 1 \in \mathbb{F}_q[x]$ is factorized into linear factors if and only if $q \equiv 1 \pmod n$.*

Remark 3.2. *The cyclotomic polynomial $\Phi_d(x) \in \mathbb{F}_q[x]$, with $\gcd(d,q) = 1$ is irreducible if and only if q is a primitive root modulus d, i.e., $\operatorname{ord}_d q = \varphi(d)$.*

Remark 3.3. *The group $\mathbb{Z}_d^*$ has a primitive root if and only if $d = 2, 4, p^i$ or $2p^i$, where p is an odd prime and $i \in \mathbb{N}^*$.*

Theorem 3.4 ([1] theorem 2.1)**.** *Let $\mathbb{F}_q$ be a finite field, and $n \in \mathbb{N}^*$ such that $q \equiv 1 \pmod n$. Then the primitive idempotents of $\frac{\mathbb{F}_q[x]}{\langle x^n-1\rangle}$ are given by*

$$e_j(x) = \frac{1}{n}\sum_{l=0}^{n-1} \zeta_n^{-jl} x^l, \quad 0 \leq j \leq n-1,$$

where $\zeta_n \in \mathbb{F}_d$ is an n-th primitive root of unity.

Theorem 3.5 ([1] theorem 3.5)**.** *Let $\mathbb{F}_q$ be a finite field and $n = p^k$, where p is an odd prime and $k \geq 1$ such that $\operatorname{ord}_n q = \varphi(n)$. Then the primitive idempotents of $\dfrac{\mathbb{F}_q[x]}{\langle x^n - 1\rangle}$ are given by*

$$e_0(x) = \frac{1}{p^k} \sum_{l=0}^{p^k-1} x^l$$

and

$$e_j(x) = \frac{1}{p^{k-j+1}} \sum_{l=0}^{p^{k-j+1}-1} x^{p^{j-1}l} - \frac{1}{p^{k-j}} \sum_{l=0}^{p^{k-j}-1} x^{p^j l} \quad 1 \leq j \leq k.$$

Observe that the representation shown here of the idempotents is the same one found by Ferraz and Polcino Milies in [3].

4. The case $p|(q-1)$

Let $\mathbb{F}_q$ be a finite field such that $p^m||(q-1)$ (i.e, $p^m|(q-1)$ and $p^{m+1} \nmid (q-1)$), where p is an odd prime and $m \geq 1$. It follows that there exists a primitive p^m-th root of unity ζ_{p^m} in $\mathbb{F}_q$. In addition, $s_j = \operatorname{ord}_{p^j} q = \begin{cases} 1 & \text{if } j \leq m \\ p^{j-m} & \text{if } j > m \end{cases}$ and, therefore, $\Phi_{p^j}(x)$ is factorized into linear factors if $j \leq m$ and, in the case $j > m$ the factorization in irreducible factors is

$$\Phi_{p^j}(x) = \Phi_{p^m}(x^{p^{j-m}}) = \prod_{\substack{l=1 \\ (p,l)=1}}^{p^m-1} (x^{p^{j-m}} - \zeta_{p^m}^l).$$

We observe that in this factorization is essential that p be an odd prime, or in the case $p = 2$ it is necessary that $q \equiv 1 \pmod 4$. Using this fact, we obtain the following new result:

Theorem 4.1. *Let $\mathbb{F}_q$ be a finite field and p is a prime such that $p^m || \gcd(p^k, q-1)$, where $m \geq 1$. If $q \equiv 1 \pmod 4$ or p is an odd prime, then the primitive idempotents of $\dfrac{\mathbb{F}_q[x]}{\langle x^{p^k} - 1\rangle}$ are of the following forms:*

(1) *For each $0 \leq j \leq p^m - 1$,*

$$e_j(x) = \frac{1}{p^k} \sum_{l=0}^{p^k-1} \zeta_{p^m}^{-jl} x^l.$$

(2) *For each $m < s \leq k$ and $0 < l < p^m$ such that $\gcd(l, p) = 1$*

$$e_{s,l}(x) = \frac{1}{p^{k+m-s}} \sum_{j=0}^{p^{k-s}-1} \zeta_{p^m}^{-lj} x^{p^{k+s-m}j}$$

Proof: Observe that

$$\frac{\mathbb{F}_q[x]}{\langle x^{p^k} - 1\rangle} \simeq \frac{\mathbb{F}_q[x]}{\langle x^{p^m} - 1\rangle} \oplus \bigoplus_{s=m+1}^{k} \frac{\mathbb{F}_q[x]}{\langle \Phi_{p^s}(x)\rangle},$$

where the second summand does not appear in the case $k = m$. Therefore, by Remark 3.1, the first case corresponds to the idempotents associated to the factor $x - \zeta_{p^m}^j \in \mathbb{F}_q[x]$ of $x^{p^k} - 1$, and by theorem 2.1 we know that the idempotent associated to this factor is of the form

$$e_j(x) = P_j(x)h_j,$$

where $P_j(x) = \frac{x^{p^m}-1}{x-\zeta_{p^m}^j}$ and $h_j \in \mathbb{F}_q$ are such that $P_j(\zeta_{p_m}^j)h_j = 1$. Observe that

$$P_j(x) = \frac{x^{p^m}-1}{x-\zeta_{p^m}^j} = \frac{x^{p^m}-(\zeta_{p^m}^j)^{p^m}}{x-\zeta_{p^m}^j} = \sum_{l=0}^{p^m-1} x^l \zeta_{p^m}^{j(p^m-1-l)} = \sum_{l=0}^{p^m-1} \zeta_n^{-jl-j} x^l,$$

and $P_j(\zeta_{p^m}^j) = p^m \zeta_{p^m}^{-j}$. Thus $h_j = \frac{\zeta_{p^m}^j}{p^m}$ and this implies the result of the first case of the theorem.

For the second case, let $x^{p^{s-m}} - \zeta_{p^m}^l$ be an irreducible factor of $\Phi_{p^s}(x)$, then the associated primitive idempotents are

$$e_{s,l} = P_{s,l}(x)h_{s,l}(x),$$

where $P_{s,l}(x) = \dfrac{x^{p^k}-1}{x^{p^{s-m}}-\zeta_{p^m}^l}$, and $h_{s,l}(x)$ is a polynomial satisfying

$$P_{s,l}(x)h_{s,l}(x) \equiv 1 \pmod{x^{p^{s-m}} - \zeta_{p^m}^l} \quad \text{and} \quad \deg(h_{s,l}(x)) < p^{s-m}.$$

Substituting $x^{p^{s-m}}$ by y, it follows that $P_{s,l}(x) = \tilde{P}(y) = \dfrac{y^{p^{k+m-s}}-1}{y-\zeta_{p^m}^l}$, and using a formal version of L'Hôpital rule, we obtain

$$\tilde{P}(y) \equiv P(1) = p^{k+m-s}\zeta_{p^m}^{l(p^{k+m-s}-1)} = p^{k+m-s}\zeta_{p^m}^{-l} \pmod{y-\zeta_{p^m}^l}$$

or, equivalently, $P_{s,l}(x) \equiv p^{k+m-s}\zeta_{p^m}^{-l} \pmod{x^{p^{s-m}} - \zeta_{p^m}^l}$.

Then $h_{s,l} = \frac{\zeta_{p^m}^l}{p^{k+m-s}}$ and

$$\begin{aligned} e_{s,l}(x) &= \frac{\zeta_{p^m}^l}{p^{k+m-s}} \frac{x^{p^k}-1}{x^{p^{s-m}}-\zeta_{p^m}^l} = \frac{\zeta_{p^m}^l}{p^{k+m-s}} \frac{x^{p^k}-(\zeta_{p^m}^l)^{p^k}}{x^{p^{s-m}}-\zeta_{p^m}^l} \\ &= \frac{1}{p^{k+m-s}} \sum_{j=0}^{p^{k-s+m}-1} \zeta_{p^m}^{l(p^{k-s+m}-j)} x^{jp^{s-m}} = \frac{1}{p^{k+m-s}} \sum_{j=0}^{p^{k-s+m}-1} \zeta_{p^m}^{-lj} x^{jp^{s-m}} \end{aligned}$$

concluding the proof. □

Remark 4.2. *The case when $p^m || (q-1)$ with $m \geq k$, is a particular case of theorem 3.4 when $n = p^k$.*

5. General Case

Let $\mathbb{F}$ be a finite field with q element and G a group p^k element, where $\gcd(q,p) = 1$. The classical method to calculate the irreducible idempotents depends of the computation of the irreducible characters $\psi : G \to \widehat{\mathbb{F}}$, where $\widehat{\mathbb{F}}$ denotes the algebraic

closure of $\mathbb{F}$, and the Galois group $Gal(\mathbb{F}(\psi),\mathbb{F})$. In fact, $e(\psi)=\frac{1}{p^k}\sum_{g\in G}\psi(g^{-1})g$ is a primitive idempotent of $\widehat{\mathbb{F}}G$ and

$$e_{\mathbb{F}}(\psi)=\sum_{\sigma\in Gal(\mathbb{F}(\psi),\mathbb{F})}\sigma\cdot e(\psi)$$

is a primitive idempotent of $\mathbb{F}G$, where σ acts on the coefficient of $e(\psi)$.

In this section, we are going to calculate the idempotent, without calculate the irreducible characters, only using the trace of some extension of $\mathbb{F}$.

Suppose that $\operatorname{ord}_p q=t>1$ and m is an integer such that $p^m||(q^t-1)$. By little Fermat theorem, it's known that $t|(p-1)$. Under such condition, $\mathbb{F}_{q^l}$ does not have a p-th primitive root of unit for all $l<t$, but $\mathbb{F}_{q^t}$ contains ζ_{p^m}, a primitive p^m-th root of unit, then $\mathbb{F}_{q^t}$ can be seen as a decomposition field of the minimal polynomial of ζ_{p^m} under $\mathbb{F}_q$, i.e., there exists an irreducible polynomial $Q(x)\in\mathbb{F}_q[x]$ of degree t, such that $Q(\zeta_{p^m})=0$.

In addition, if $\tau\in Gal(\mathbb{F}_{q^t},\mathbb{F}_q)$ is the Fröbenius automorphism $a\overset{\tau}{\longmapsto}a^q$, then

$$\{\tau^j(\zeta_{p^m})|j=0,1,\ldots,t-1\}=\{\zeta_{p^m}^{q^j}|j=0,1,\ldots,t-1\}$$

is the set of conjugates of ζ_{p^m} over $\mathbb{F}_q$. In general, for all $a\in\mathbb{F}_{q^t}$, the function

$$\begin{array}{rccl}\sigma_1: & \mathbb{F}_{q^t} & \to & \mathbb{F}_q\\ & a & \mapsto & a+\tau(a)+\tau^2(a)+\cdots+\tau^{(t-1)}(a),\end{array}$$

is well defined.

By the previous section, we show the explicit form of the primitive idempotents of $\dfrac{\mathbb{F}_{q^t}[x]}{\langle x^{p^k}-1\rangle}$. The next theorem uses this representation in order to calculate the form of the primitive idempotents of $\dfrac{\mathbb{F}_q[x]}{\langle x^{p^k}-1\rangle}$.

Theorem 5.1. *Let $\mathbb{F}_q$ be a finite field and assume p is an odd prime such that $\operatorname{ord}_p q=t>1$ and $p^m||(q^t-1)$, where $m\geq 1$. Then the primitive idempotents of $\dfrac{\mathbb{F}_q[x]}{x^{p^k}-1}$ are of the following forms:*

(1) $e_0(x)=\dfrac{1}{p^k}\displaystyle\sum_{l=0}^{p^k-1}x^l$.

(2) *For each $0<j\leq p^m-1$,*

$$e_j(x)=\frac{1}{p^k}\sum_{l=0}^{p^k-1}\sigma_1(\zeta_{p^m}^{-jl})x^l,$$

where e_i and e_j are the same idempotents if and only if $i\equiv jq^u \pmod{p^m}$ for some $u\in\mathbb{Z}$, and, therefore, there are $\frac{p^m-1}{t}$ different primitive idempotents of this type.

(3) *For each $m<s\leq k$ and $0<l<p^m$ such that $\gcd(l,p)=1$,*

$$e_{s,l}(x)=\frac{1}{p^{k+m-s}}\sum_{j=0}^{p^{k-s+m}-1}\sigma_1\left(\zeta_{p^m}^{-lj}\right)x^{p^{s-m}j},$$

where e_{s,l_1} and e_{s,l_2} are the same idempotents if and only if $l_1 \equiv l_2 q^u \pmod{p^m}$, for some $u \in \mathbb{N}$, and, therefore, for each s fixed, there are $\frac{\varphi(p^{k-s+m})}{t}$ different primitive idempotents of this type.

Remark 5.2. *In* [8]*, using a cyclotomic cosets method, was studied the particular case when $m = 1$.*

Proof: Let $E(x)$ be a primitive idempotent of $\dfrac{\mathbb{F}_q[x]}{\langle x^{p^k}-1\rangle}$. It follows that $E(x)$ is also an idempotent of $\dfrac{\mathbb{F}_{q^t}[x]}{\langle x^{p^k}-1\rangle}$ and therefore $E(x)$ is a direct sum of primitive idempotents of $\dfrac{\mathbb{F}_{q^t}[x]}{\langle x^{p^k}-1\rangle}$. In the case that $E(x)$ be also primitive in $\dfrac{\mathbb{F}_{q^t}[x]}{\langle x^{p^k}-1\rangle}$, then by theorem 4.1 we know the unique idempotent with this propriety is $e_0(x)$, i.e., the case when $j = 0$ and therefore $E(x) = e_0(x)$.

Now, suppose that $E(x) \neq e_0(x)$, and let $e(x) \in \dfrac{\mathbb{F}_{q^t}[x]}{\langle x^{p^k}-1\rangle}$ be a primitive idempotent such that $e(x) \cdot E(x) = e(x) \notin \dfrac{\mathbb{F}_q[x]}{\langle x^{p^k}-1\rangle}$. Since $\tau(E(x)) = E(x)$, it follows that $\tau(e(x))$ is also a direct summand of $E(x)$. In addition, it is known that $\tau^r(\zeta_{p^m}^j) = \zeta_{p^m}^j$ if and only if $j(q^r-1)$ is divisible by p^m and this is equivalent to $p^m | j$ or $t | r$. From this, we conclude that $\tau^r(e(x)) = e(x)$ if and only if $t|r$, thus

$$\{e(x), \tau(e(x)), \ldots, \tau^{(t-1)}(e(x))\}$$

is a list of different idempotents that are direct summands of $E(x)$. Finally, since $\sigma_1(e(x)) := e(x) + \tau(e(x)) + \cdots + \tau^{(t-1)}(e(x)) \in \dfrac{\mathbb{F}_q[x]}{\langle x^{p^k}-1\rangle}$ and using the fact that $E(x)$ is primitive, we conclude that $E(x) = \sigma_1(e(x))$, in other words, we can obtain every primitive idempotent of $\dfrac{\mathbb{F}_q[x]}{\langle x^{p^k}-1\rangle}$ different of $e_0(x)$ from the idempotent of $\dfrac{\mathbb{F}_{q^t}[x]}{\langle x^{p^k}-1\rangle}$ and the ring homomorphism σ_1. Thus, the other cases of the theorem follow directly from the cases (1) and (2) of Theorem 4.1. □

6. Sage implementation and examples

In this section, some examples are shown explicitly. In order to find these idempotents, we have implemented the last theorem in the SAGE program[1], as it is shown in the following code:

First we defined the field $\mathbb{F}_{q^t}$, the p^m-th root of the unity, and the polynomial ring $\mathbb{F}_{q^t}[x]$

```
sage: k.<a>=GF(q^t,'a');
sage: b=a^((q^t-1)/p^m);
sage: F.<x>=PolynomialRing(k,'x')
```

Implementation the function $\sigma_1(\zeta_{p^m}^{lu})$

```
sage: def sigma(l,u):
...       sumconj=sum([b^(l*u*q^i) for i in range(0,t)]);
...       return(sumconj)
```

[1] `http://www.sagemath.org`

The idempotents of the second type

```
sage: def Idemp2(l):
...       v = []
...       for i in range(0,p^k):
...          v.append( sigma(q,t,l,i)/p^k)
...       Poli= sum([v[j]*x^j for j in range(0,p^k)])
...       return Poli
```

The idempotents of the third type

```
sage: def Idemp3(l,s):
...       v = []
...       for j in range(0,p^(k-s+m)):
...          v.append( sigma(q,t,l,p^(k-s+m)-j)/p^(k+m-s))
...       Poli= sum([v[j]*x^(p^(s-m)*j) for j in range(0,p^(k-s+m))])
...       return Poli
```

Example 6.1. *In the group ring* $\mathbb{F}_{17}C_{13^2}$*, since* $13||17^6-1$*, then* $t=6$ *and* $m=1$*. Thus, there exists* $\frac{13-1}{6}=2$ *primitive idempotents of the second type, and making* $s=2=k$*, there exists* $\frac{\varphi(13)}{6}=2$ *primitive idempotents of the third type. In fact, the primitive idempotents are:*

- $e_0 = 16\sum_{i=0}^{168} x^i$;
- *two idempotents of the second type*

$$e_1 = \sum_{i=0}^{12} x^{13i}(5x^{12}+13x^{11}+5x^{10}+5x^9+13x^8+13x^7+13x^6+ \\ +13x^5+5x^4+5x^3+13x^2+5x+11)$$

$$e_2 = \sum_{i=0}^{12} x^{13i}(13x^{12}+5x^{11}+13x^{10}+13x^9+5x^8+5x^7+5x^6+ \\ +5x^5+13x^4+13x^3+5x^2+13x+11)$$

- *and two idempotents of the third type*

$$e_{2,1} = 14x^{156}+16x^{143}+14x^{130}+14x^{117}+16x^{104}+16x^{91}+16x^{78}+ \\ +16x^{65}+14x^{52}+14x^{39}+16x^{26}+14x^{13}+7$$

$$e_{2,2} = 16x^{156}+14x^{143}+16x^{130}+16x^{117}+14x^{104}+14x^{91}+14x^{78}+ \\ +14x^{65}+16x^{52}+16x^{39}+14x^{26}+16x^{13}+7$$

References

[1] Arora, S. K., Pruthi M. *Minimal Codes of Prime-Power Length*. Finite Fields and Their Aplications **3** (1997) 99-113

[2] Arora, S. K., Pruthi M. *Minimal Cyclic Codes of Length* $2p^n$. Finite Fields and Their Aplications **5** (1999) 177-187

[3] Ferraz, R., Polcino Milies C.. *Idempotents in group algebras and minimal abelian codes*. Finite Fields and Their Applications **13** (2007) 382-393

[4] Curtis, C.W., Reiner, I., *Representation Theory of finite Groups and associative Algebra*. Interscience Publishers 1962

[5] Lidl, R., Niederreiter, H. *Finite Fields*. Encyclopedia of Mathematics and Its Applications, Vol 20, Addison-Wesley 1983.

[6] Fitzgerald R.W., Yucas J.L.:*Explicit factorization of cyclotomic and Dickson polynomials over finite fields*. Arithmetic of Finite Fields. Lecture Notes in Computer Science, vol. 4547, pp. 1-10. Springer, Berlin (2007).

[7] Meyn H. *Factorization of the cyclotomic polynomials* $x^{2^n}+1$ *over finite fields*. Finite Fields Appl. 2, (1996) 439-442

[8] Sharma, A., Bakshi, G., Dumir, V. C., Raka, M., *Cyclotomic numbers and primitive idempotents in the ring* $GF(q)[x]/(x^{p^n}-1)$. Finite Fields and Their Aplications **10** (2004), 653-673.

Departamento de Matemática, Universidade Federal de Minas Gerais, UFMG, Belo Horizonte, MG, 30123-970, Brazil,

E-mail address: fbrocher@mat.ufmg.br

E-mail address: carmita@mat.ufmg.br